

# ***The Sorption of Selected Radionuclides in Sedimentary Interbed Soils from the Snake River Plain***

*June 2001*



*Idaho National Engineering and Environmental Laboratory  
Bechtel BWXT Idaho, LLC*

# **The Sorption of Selected Radionuclides in Sedimentary Interbed Soils from the Snake River Plain**

**Christopher J. Grossman  
Robert A. Fjeld  
John T. Coates  
Alan W. Elzerman**

**Department of Environmental Engineering and Science  
Clemson University  
Clemson, South Carolina**

**June 2001**

**Idaho National Engineering and Environmental Laboratory  
Environmental Restoration Project  
Idaho Falls, Idaho 83415**

**Prepared under Subcontract No. K99-181044  
for the  
U.S. Department of Energy  
Assistant Secretary for Environmental Management  
Under DOE Idaho Operations Office  
Contract DE-AC07-99ID13727**

Note: This document was produced for the Idaho National Engineering and Environmental Laboratory (INEEL) by an external subcontractor, and may not have been technically reviewed or edited to meet the INEEL publications standard.

**TASK 7 FINAL REPORT**

**THE SORPTION OF SELECTED RADIONUCLIDES IN SEDIMENTARY INTERBED  
SOILS FROM THE SNAKE RIVER PLAIN**

**Submitted by**

**Christopher J. Grossman, Robert A. Fjeld, John T. Coates, and Alan W. Elzerman  
Department of Environmental Engineering and Science  
Clemson University**

**to**

**Bechtel BWXT Idaho  
CONTRACT # K99-181044**

**June 27, 2001**

**Technical Contact:**

**John T. Coates  
864-656-1540 (voice)  
864-656-0672 (fax)  
e-mail: [coates@clemson.edu](mailto:coates@clemson.edu)**

## ABSTRACT

Batch kinetic and equilibrium sorption experiments were conducted to characterize the sorption behavior of a selected suite of contaminants (uranium, neptunium, americium, and chromium), which were buried in the Radioactive Waste Management Complex at the Idaho National Engineering and Environmental Laboratory. The principal objective of this research was to characterize the sorption isotherms for the selected radionuclides under conditions representative of actual field conditions beneath the Radioactive Waste Management Complex and to examine the variability of their sorption behaviors among soils collected at different depths. A secondary objective was to examine the effect of bicarbonate, sulfate, and fluoride, which are common groundwater ligands, on sorption behavior.

Batch experiments were conducted with depth-discrete core samples of sedimentary interbed from beneath the Radioactive Waste Management Complex. Sorbed phase concentrations were inferred from changes in aqueous phase concentrations. The aqueous phase concentration was operationally defined as the remainder after filtration at nominal 12 nm measured by liquid scintillation counting.

Sorption behaviors for  $^{233}\text{U}$  and  $^{237}\text{Np}$  were nonlinear. The Freundlich and Langmuir sorption models were fit to the equilibrium data. The Freundlich model fit the data better than the Langmuir model for both  $^{233}\text{U}$  and  $^{237}\text{Np}$ . Spatial variation of sorption behavior was found to be small for both contaminants.

Carbonate was found to have a strong influence on  $^{233}\text{U}$  behavior and masked any effects from sulfate and fluoride. Carbonate, sulfate, and fluoride had no effect on  $^{237}\text{Np}$  sorption behavior. The predominant sorbing neptunyl species was predicted to be  $\text{NpO}_2^+$ .

Large losses of  $^{51}\text{Cr}$  and  $^{241}\text{Am}$  in blank samples made the determination of their sorption behavior problematic. For  $^{51}\text{Cr}$ , the losses were likely due to

The loss of aqueous phase  $^{51}\text{Cr}$  was lowest in samples with the highest solids concentration. Additionally, large losses of  $^{51}\text{Cr}$  were observed in equilibrium blanks. Apparent distribution coefficients were determined for the samples with the highest solids concentrations to minimize the effect of precipitation and estimate an upper limit of sorption affinity between  $^{51}\text{Cr}$  and the soils. The apparent distribution coefficients were small suggesting that the sorption affinity of  $^{51}\text{Cr}$  for these soils was small.

Large losses of  $^{241}\text{Am}$  were observed in short-term blanks due to formation of insoluble americium carbonates. Consequently, equilibrium studies were not conducted.

## TABLE OF CONTENTS

	Page
TITLE PAGE.....	i
ABSTRACT .....	ii
LIST OF TABLES .....	vi
LIST OF FIGURES .....	ix
CHAPTER	
1 INTRODUCTION .....	1
2 OBJECTIVES.....	5
3 ENVIRONMENTAL GEOCHEMISTRY .....	6
Aqueous Contaminant Speciation.....	6
Chromium .....	7
Uranium .....	8
Neptunium .....	9
Americium .....	11
Sorption.....	12
Interfacial Forces .....	12
Surface Complex Formation .....	13
Sorption Models .....	14
4 MATERIALS AND METHODOLOGY .....	17
Materials .....	17
Groundwater Simulant .....	17
Sedimentary Interbed.....	19
Radionuclide Preparations.....	20
Methodology .....	22
Batch Kinetic Studies .....	22
Batch Equilibrium Studies .....	23
Inorganic Carbon Analysis .....	24

## Table of Contents (Continued)

	Page
5 RESULTS AND DISCUSSION .....	25
Batch Kinetic Studies .....	25
Uranium .....	25
Neptunium .....	27
Americium .....	29
Chromium .....	31
Batch Equilibrium Studies .....	34
Uranium .....	34
Neptunium .....	40
Chromium .....	44
Soil Property Correlations .....	48
Ligand Studies .....	50
Uranium .....	52
Neptunium .....	55
6 CONCLUSIONS.....	61
Batch Equilibrium Studies .....	61
Environmental Significance.....	63
Recommendations .....	65
APPENDICES .....	66
A Soils Characterization Data.....	67
B Experimental Data .....	75
C Uranium Isotherms.....	88
D Neptunium Isotherms.....	102
REFERENCES.....	116



## LIST OF TABLES

Table		Page
Table 4.1	Composition of Snake River Plain Aquifer (SRPA) groundwater and the groundwater simulants used in this study. ....	18
Table 4.2	Summary of bulk characteristics for INEEL soils where available.....	20
Table 4.3	Summary of clay fraction characteristics for INEEL soils where available.....	21
Table 4.4	Summary of mineral oxide characteristics for INEEL soils where available. ....	21
Table 5.1	Summary of kinetic study results for uranium on soils 7DS01701KD and I1S-INEEL-109. ....	25
Table 5.2	Summary of kinetic study results for neptunium on soils 7DS01701KD and I1S-INEEL-109. ....	27
Table 5.3	Summary of americium losses from short-term blanks (no soil) in GWS. ....	29
Table 5.4	Summary of chromium losses from short-term blanks (no soil) in GWS. ....	31
Table 5.5	Summary of kinetic study results for chromium on soils 7DS00901KD and I4D-INEEL-234.....	34
Table 5.6	Summary of sorption models with respective statistics of model fit determined from the results of the equilibrium study for uranium on all soils. ....	38
Table 5.7	Comparison of uranium Freundlich isotherm parameters for soils from Oak Ridge National Laboratory (ORNL), Savannah River Site (SRS), Hanford Reservation (HR), and the Idaho National Engineering and Environmental Laboratory (INEEL). ....	39
Table 5.8	Summary of sorption models with respective statistics of model fit determined from the results of the equilibrium study for neptunium on all soils. ....	43

## List of Tables (Continued)

Table	Page
Table 5.9 Summary of equilibrium study data for chromium on fourteen soil samples and blanks. ....	45
Table 5.10 Summary of apparent distribution ratios from samples with solids concentrations that approximated 50,000 mg-L <sup>-1</sup> for all soils. ....	48
Table 5.11 Summary of linear sorption model with respective statistics of model fit determined from the results of the equilibrium study for uranium at the three lowest aqueous phase concentrations on soils for which characterization data were available. ....	49
Table 5.12 Summary of linear sorption model with respective statistics of model fit determined from the results of the equilibrium study for neptunium at the three lowest aqueous phase concentrations on soils for which characterization data were available. ....	49
Table 5.13 Summary of statistical correlations between the fitted linear sorption model and available soil properties. ....	50
Table 5.14 Summary of inorganic carbon concentrations determined in background (no activity) samples at equilibrium for soils I2S-INEEL-105 and I4D-INEEL-234. ....	51
Table 5.15 Summary of Freundlich parameters with statistics of fit determined from ligand study data for neptunium on soil I2S-INEEL-105 and I4D-INEEL-234. ....	58
Table A.1 Fraction of each soil passing #5 (Dia. < 2 mm) sieve. ....	74
Table B.1 Summary of experimental solids concentrations for chromium kinetic and equilibrium studies. ....	76
Table B.2 Summary of experimental solids concentrations for uranium kinetic and equilibrium studies. ....	77
Table B.3 Summary of experimental solids concentrations for neptunium kinetic and equilibrium studies. ....	78
Table B.4 Summary of experimental initial aqueous phase concentrations for radionuclide kinetic and equilibrium studies. ....	79

## List of Tables (Continued)

Table	Page
Table B.5 Summary of initial uranium-233 aqueous activity and mass concentrations and equivalent natural uranium activity and mass concentrations. ....	80
Table B.6 Summary of equilibrium study data for uranium on fourteen soil samples and blanks in GWS. ....	81
Table B.7 Summary of ligand study data for uranium on two soil samples and blanks in GWS(-CO <sub>3</sub> <sup>2-</sup> ) and GWS(-CO <sub>3</sub> <sup>2-</sup> , SO <sub>4</sub> <sup>2-</sup> , F <sup>-</sup> ) .....	83
Table B.8 Summary of equilibrium study data for neptunium on fourteen soil samples and blanks in GWS. ....	84
Table B.9 Summary of ligand study data for neptunium on two soil samples and blanks in GWS(-CO <sub>3</sub> <sup>2-</sup> ) and GWS(-CO <sub>3</sub> <sup>2-</sup> , SO <sub>4</sub> <sup>2-</sup> , F <sup>-</sup> ) ...	86

## LIST OF FIGURES

Figure	Page
Figure 3.1 Adsorption mechanisms at oxide surfaces.....	14
Figure 5.1 Results of kinetic study for uranium on two soils: I1S-INEEL-109 (■); 7DS01701KD (●).....	26
Figure 5.2 Results of kinetic study for neptunium on two soils: I1S-INEEL-109 (■); 7DS01701KD (●).....	28
Figure 5.3 Results of kinetic study for chromium on two soils: I4D-INEEL-234 (■); 7DS00901KD (●).....	33
Figure 5.4 Results of equilibrium study for uranium on soil: 7DS01701KD (◇).....	35
Figure 5.5 Application of sorption models to results of equilibrium study for uranium on soil 7DS01701KD (◇). Solid line represents the fitted Freundlich model. Dashed line represents the fitted Langmuir model. ....	37
Figure 5.6 Results of equilibrium study for neptunium on soil: 7DS01701KD (◇).....	41
Figure 5.7 Application of sorption models to results of equilibrium study for neptunium on soil 7DS01701KD (◇). Solid line represents the fitted Freundlich model. Dashed line represents the fitted Langmuir model. ....	42
Figure 5.8 Results of ligand study for uranium on soil I2S-INEEL-105 for GWS (◇); GWS (-CO <sub>3</sub> <sup>2-</sup> ) (□); GWS (-CO <sub>3</sub> <sup>2-</sup> , SO <sub>4</sub> <sup>2-</sup> , F <sup>-</sup> ) (Δ).....	53
Figure 5.9 Results of ligand study for uranium on soil I4D-INEEL-234 for GWS (◇); GWS (-CO <sub>3</sub> <sup>2-</sup> ) (□);GWS (-CO <sub>3</sub> <sup>2-</sup> , SO <sub>4</sub> <sup>2-</sup> , F <sup>-</sup> ) (Δ). ....	54
Figure 5.10 Results of ligand study for neptunium on soil I2S-INEEL-105 for GWS (◇); GWS (-CO <sub>3</sub> <sup>2-</sup> ) (□); GWS (-CO <sub>3</sub> <sup>2-</sup> , SO <sub>4</sub> <sup>2-</sup> , F <sup>-</sup> ) (Δ).....	56
Figure 5.11 Results of ligand study for neptunium on soil I4D-INEEL-234 for GWS (◇); GWS (-CO <sub>3</sub> <sup>2-</sup> ) (□);GWS (-CO <sub>3</sub> <sup>2-</sup> , SO <sub>4</sub> <sup>2-</sup> , F <sup>-</sup> ) (Δ).....	57
Figure 5.12 Computed speciation of NpO <sub>2</sub> <sup>+</sup> as a function of pH in GWS via MINTEQA2 (Allison <i>et al.</i> , 1991).....	60

## List of Figures (Continued)

Figure	Page
Figure A.1 Particle size distribution for soil 7DS00101KD. Work performed by the SwRI under INEEL ER-SOW-371. ....	68
Figure A.2 Particle size distribution for soil 7DS00301KD. Work performed by the SwRI under INEEL ER-SOW-371. ....	69
Figure A.3 Particle size distribution for soil 7DS00501KD. Work performed by the SwRI under INEEL ER-SOW-371. ....	70
Figure A.4 Particle size distribution for soil 7DS00701KD. Work performed by the SwRI under INEEL ER-SOW-371. ....	71
Figure A.5 Particle size distribution for soil 7DS00901KD. Work performed by the SwRI under INEEL ER-SOW-371. ....	72
Figure A.6 Particle size distribution for soil 7DS02301KD. Work performed by the SwRI under INEEL ER-SOW-371. ....	73
Figure B.1 Calibration curve for inorganic carbon determinations. ....	87
Figure C.1 Application of sorption models to results of equilibrium study for uranium on soil 7DS00101KD (◇). Solid line represents the fitted Freundlich model. Dashed line represents the fitted Langmuir model. ....	89
Figure C.2 Application of sorption models to results of equilibrium study for uranium on soil 7DS00301KD (◇). Solid line represents the fitted Freundlich model. Dashed line represents the fitted Langmuir model. ....	90
Figure C.3 Application of sorption models to results of equilibrium study for uranium on soil 7DS00501KD (◇). Solid line represents the fitted Freundlich model. Dashed line represents the fitted Langmuir model. ....	91
Figure C.4 Application of sorption models to results of equilibrium study for uranium on soil 7DS00701KD (◇). Solid line represents the fitted Freundlich model. Dashed line represents the fitted Langmuir model. ....	92
Figure C.5 Application of sorption models to results of equilibrium study for uranium on soil 7DS00901KD (◇). Solid line represents the fitted Freundlich model. Dashed line represents the fitted Langmuir model. ....	93

## List of Figures (Continued)

Figure	Page
Figure C.6 Application of sorption models to results of equilibrium study for uranium on soil 7DS02301KD ( $\diamond$ ). Solid line represents the fitted Freundlich model. Dashed line represents the fitted Langmuir model. ....	94
Figure C.7 Application of sorption models to results of equilibrium study for uranium on soil I2S-INEEL-105 ( $\diamond$ ). Solid line represents the fitted Freundlich model. Dashed line represents the fitted Langmuir model. ....	95
Figure C.8 Application of sorption models to results of equilibrium study for uranium on soil I1S-INEEL-109 ( $\diamond$ ). Solid line represents the fitted Freundlich model. Dashed line represents the fitted Langmuir model. ....	96
Figure C.9 Application of sorption models to results of equilibrium study for uranium on soil I4D-INEEL-224 ( $\diamond$ ). Solid line represents the fitted Freundlich model. Dashed line represents the fitted Langmuir model. ....	97
Figure C.10 Application of sorption models to results of equilibrium study for uranium on soil I3D-INEEL-229 ( $\diamond$ ). Solid line represents the fitted Freundlich model. Dashed line represents the fitted Langmuir model. ....	98
Figure C.11 Application of sorption models to results of equilibrium study for uranium on soil I4D-INEEL-231 ( $\diamond$ ). Solid line represents the fitted Freundlich model. Dashed line represents the fitted Langmuir model. ....	99
Figure C.12 Application of sorption models to results of equilibrium study for uranium on soil I1D-INEEL-234 ( $\diamond$ ). Solid line represents the fitted Freundlich model. Dashed line represents the fitted Langmuir model. ....	100
Figure C.13 Application of sorption models to results of equilibrium study for uranium on soil I4D-INEEL-234 ( $\diamond$ ). Solid line represents the fitted Freundlich model. Dashed line represents the fitted Langmuir model. ....	101

## List of Figures (Continued)

Figure	Page
Figure D.1 Application of sorption models to results of equilibrium study for neptunium on soil 7DS00101KD ( $\diamond$ ). Solid line represents the fitted Freundlich model. Dashed line represents the fitted Langmuir model. ....	103
Figure D.2 Application of sorption models to results of equilibrium study for neptunium on soil 7DS00301KD ( $\diamond$ ). Solid line represents the fitted Freundlich model. Dashed line represents the fitted Langmuir model. ....	104
Figure D.3 Application of sorption models to results of equilibrium study for neptunium on soil 7DS00501KD ( $\diamond$ ). Solid line represents the fitted Freundlich model. Dashed line represents the fitted Langmuir model. ....	105
Figure D.4 Application of sorption models to results of equilibrium study for neptunium on soil 7DS00701KD ( $\diamond$ ). Solid line represents the fitted Freundlich model. Dashed line represents the fitted Langmuir model. ....	106
Figure D.5 Application of sorption models to results of equilibrium study for neptunium on soil 7DS00901KD ( $\diamond$ ). Solid line represents the fitted Freundlich model. Dashed line represents the fitted Langmuir model. ....	107
Figure D.6 Application of sorption models to results of equilibrium study for neptunium on soil 7DS02301KD ( $\diamond$ ). Solid line represents the fitted Freundlich model. Dashed line represents the fitted Langmuir model. ....	108
Figure D.7 Application of sorption models to results of equilibrium study for neptunium on soil I2S-INEEL-105 ( $\diamond$ ). Solid line represents the fitted Freundlich model. Dashed line represents the fitted Langmuir model. ....	109
Figure D.8 Application of sorption models to results of equilibrium study for neptunium on soil I1S-INEEL-109 ( $\diamond$ ). Solid line represents the fitted Freundlich model. Dashed line represents the fitted Langmuir model. ....	110

## List of Figures (Continued)

Figure	Page
Figure D.9 Application of sorption models to results of equilibrium study for neptunium on soil I4D-INEEL-224 ( $\diamond$ ). Solid line represents the fitted Freundlich model. Dashed line represents the fitted Langmuir model. ....	111
Figure D.10 Application of sorption models to results of equilibrium study for neptunium on soil I3D-INEEL-229 ( $\diamond$ ). Solid line represents the fitted Freundlich model. Dashed line represents the fitted Langmuir model. ....	112
Figure D.11 Application of sorption models to results of equilibrium study for neptunium on soil I4D-INEEL-231 ( $\diamond$ ). Solid line represents the fitted Freundlich model. Dashed line represents the fitted Langmuir model. ....	113
Figure D.12 Application of sorption models to results of equilibrium study for neptunium on soil I1D-INEEL-234 ( $\diamond$ ). Solid line represents the fitted Freundlich model. Dashed line represents the fitted Langmuir model. ....	114
Figure D.13 Application of sorption models to results of equilibrium study for neptunium on soil I4D-INEEL-234 ( $\diamond$ ). Solid line represents the fitted Freundlich model. Dashed line represents the fitted Langmuir model. ....	115